Setting the **Global** Standard for Quality in Laboratory Testing 
 P: +1.610.688.0100
 950 West V

 Toll Free (US): 877.447.1888
 Suite 2500

 F: +1.610.688.0700
 Wayne, PA :

 E: customerservice@clsi.org
 Suite 2500

950 West Valley Road Suite 2500 Wayne, PA 19087 USA



## 22 October 2019

To: Recipients of VET06, 1st ed.

From: Jennifer K. Adams, MT(ASCP), MSHA Vice President, Standards and Quality

Subject: Correction

This notice is intended to inform users of corrections made to CLSI document VET06, *Methods for Antimicrobial Susceptibility Testing of Infrequently Isolated or Fastidious Bacteria Isolated From Animals*, 1st ed. The corrections are described below and shown as highlighted and/or stricken text in the table excerpt.

## Table 17. Gallibacterium anatis-Information and Breakpoints for Broth Microdilution Susceptibility Testing:

The minimal inhibitory concentration (MIC) intermediate ranges are listed incorrectly for penicillin, clindamycin, and erythromycin. The MIC intermediate range for penicillin has been corrected to read "-," the MIC intermediate range for clindamycin has been corrected to read "1-2," and the MIC intermediate range for erythromycin has been corrected to read "1-4."

Table 17.	Gallibacterium	anatis-Information	and	Breakpoints	for	Broth	Microdilution
Susceptibility Testing							

Antimicrobial		MIC (µg/mL) Interpretive Categories and Breakpoints							
Class	Antimicrobial Agent	S	I	R	Comments				
B-LACTAMS									
	Penicillin	≤0.125	<mark>1-2</mark> -	≥0.25					
LINCOSAMIDE									
	Clindamycin	≤0.5	<mark>— 1-2</mark>	≥4					
MACROLIDES			•	•					
	Erythromycin	≤0.5	<mark>0.5</mark> 1-4	≥8					

Abbreviations: I, intermediate; MIC, minimal inhibitory concentration; R, resistant; S, susceptible.

If you require any additional clarification regarding these corrections, please contact CLSI Customer Service (<u>customerservice@clsi.org</u>).

We appreciate your commitment to CLSI and regret any inconvenience.